

22-0-5

PCSCU

GUARANTEED ANALYSIS

Total Nitrogen (N)	22.0%
22.0% Urea nitrogen*	
Soluble Potash (K ₂ O)	5.0%
Sulfur (S)	0.6%
Iron (Fe)	1.0%

Primary Plant foods derived from: Urea, Muriate of Potash and Poly Coated Sulfur Coated Urea. Sulfur derived from Poly Coated Sulfur Coated Urea. Iron derived from Iron Oxy-Sulfate.

*6.6% slowly available urea nitrogen from Poy Coated Sulfur Coated Urea. 544-3030 polyocket

Net Wt. 50 lbs. (22.7 kg)

ACTIVE INGREDIENTS

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	1.040%
Dimethylamine salt of (+)-(R)-2-[2-methyl-4-chlorophenoxy] propionic acid	0.230%
Dimethylamine salt of dicamba [3,6-dichloro-o-anisic acid]	0.108%
INERT INGREDIENTS	98.622%
TOTAL INGREDIENTS	100.000%

EPA REG. NO. 2217-561-57131

EPA EST. NO. 57131-WI-1

Controls chickweed, dandelion, knotweed, plantain, henbit, spurge and many other weeds as listed.



THIS PRODUCT CONTAINS:

- 17.20 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.86%
- 3.80 lbs. (+)-(R)-2-[2-methyl-4-chlorophenoxy] propionic acid equivalent per ton or 0.19%
- 1.80 lbs. 3,6-dichloro-o-anisic acid equivalent per ton or 0.09%

Isomer Specific by AOAC Methods

TRIMEC® is a registered trademark of FBI/GORDON CORPORATION

Contains the single isomer form of Mecoprop-p.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional Precautionary Statements.



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Avoid inhaling dust or contact with skin, eyes and clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

FIRST AID

- If in eyes:** • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.
- If on skin or clothing:** • Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 - 20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Most cases of groundwater contamination involving phenoxy herbicides such as MCPP-p and 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCPP-p and 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

BROADLEAF WEED CONTROL

CONTAINS TRIMEC® HERBICIDE

DIRECTIONS FOR USE

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: For Paper Bags, Plastic Bags, and Nonrefillable Bulk Bags - Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or if allowed by State and local authorities by burning. If burned stay out of smoke.

WHERE TO USE

This product is a Trimec® herbicide containing fertilizer, 2,4-D, mecoprop-p, and dicamba. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash should be applied according to soil tests or State Extension Service recommendations. This product is intended to be applied by commercial applicators, noncommercial users, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated. Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; for example: hospitals, nursing homes, schools, museums, libraries, sport facilities (including golf courses), and office buildings may be treated. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes. This product may be applied to newly seeded cool season turfgrass. Applications are recommended only after the third mowing or after the grass seedlings have tilled and have developed an adequate secondary root system. This product may be applied to newly established warm season turfgrass. Treatments should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations, and low application rates (1/2 of the recommendation for established turfgrass) are suggested. A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals. Do not use this product on St. Augustinegrass, dichondra, carpetgrass, closely-mowed bentgrass putting greens, or turf containing desirable clovers.

HOW MUCH TO APPLY

For Kentucky bluegrass, fescue spp., perennial ryegrass, bermudagrass, and zoysiagrass: Apply 130-174 pounds of this product per acre of established turfgrass. Or the equivalent application rates are 3.0-4.0 pounds of product per 1,000 square feet of turfgrass. The maximum application rate to turf is 1.5 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year. Sequential or follow-up applications may be required for dense infestations of perennial and biennial weeds. Also, adverse environmental conditions prior or after application may reduce the effectiveness. If required, the treatment interval between the first and second applications should be approximately 30 days. Drop spreaders, rotary spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer. Refer to the rate table for the appropriate bag sizes and the area to be treated.

APPLICATION SUGGESTIONS:

1. Spring and fall applications are recommended for the best control. However, this product can be applied any time broadleaf weeds are present and actively growing.
2. Mow the turfgrass 1 to 2 days before application.
3. Water the turfgrass thoroughly (> 1/2 inch) at least 1 to 2 days before application.
4. Schedule applications in the morning when dew is present. Or sprinkle or irrigate lightly (<1/4 inch) before application to provide maximum control.
5. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.



PROFESSIONAL TURF FERTILIZER

4 lbs Per 1,000

SPREADERS	Sq. Ft.
Lesco Rotary	F-G
Spyker 24-22	4
Scott's R8A	1
Prizelawn CB Rotary	4
Earthway 2400 Rotary	17
Lely Broadcast	3.4

CALIBRATION INSTRUCTIONS:

To provide proper distribution, calibrate fertilizer spreader before application. For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

DISTRIBUTION AND SPREADER PATTERNS: Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

COVERAGE: To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to 12,500 square feet.

SINGLE PASS DISTRIBUTION: First spread the borders of your lawn. Then spread the center by spreading in straight line passes. Spread at a uniform walking speed to insure even coverage. To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes.

Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.

WEEDS CONTROLLED:

Dandelion, chickweed, plantain, and clover are controlled effectively with this product. A partial list of susceptible weeds is shown below.

Bedstraw	Dock	Mallow	Poison Oak	Thistle
Black medic	Ground Ivy	Morningglory	Purslane	Wild Carrot
Buckhorn	Healall	Oxalis	Ragweed	Wild Garlic
Burdock	Herbit	Peppergrass	Sheep sorrel	Wild Lettuce
Chickweed	Knotweed	Pigweed	Sheepspurge	Wild Onion
Clover	Lambsquarters	Plantain	Speedwell	Yarrow
Dandelion	Lespedeza	Poison Ivy	Spurge	

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.



Distributed By Eau Claire Co-op Oil Co., P.O. Box 837, Eau Claire, WI 54702